

DIRECT™



DRIFT RETARDANT/DEPOSITION AID

PRINCIPAL FUNCTIONING AGENTS:

Polyvinyl Polymer (Polyacrylamide).....	30%
Inert Ingredients (Formulating Aid)	70%
TOTAL.....	100%

All ingredients are approved for use under CFR 40,180.1001(c)

CAUTION: KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

Before using this product, read the entire label, including the conditions of sale. May be harmful if swallowed, inhaled, or contact is made with skin or eyes. Breathing excessive concentration of vapors can cause dizziness. Always wear personal protective equipment as mandated by the label of the pesticide tank mix partner(s) being used. Follow manufacturer's instructions for cleaning and maintaining PPE.

DIRECT will retard, but cannot totally eliminate, drift hazards due to variations in pesticide usage, application conditions and techniques, as well as sensitive vegetation near the target application area. Minimizing spray drift is the responsibility of the applicator.

STATEMENT OF PRACTICAL TREATMENT

- If Swallowed:** Seek medical attention immediately.
- If In Eyes:** Flush with water for 15 minutes. Seek medical attention immediately.
- If On Skin:** Wash skin contact area with plenty of soap and water for 15 minutes. Seek medical attention if irritation persists.
- If Inhaled:** Remove to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen.

GENERAL INFORMATION

DIRECT Drift Retardant and Deposition Aid reduces drift risk and maximizes pesticide performance by binding Ultra Small Spray Particles together into larger droplets which are more capable of hitting their intended target. DIRECT's refined use rates and low "phase out" formulation make it the easiest to use liquid drift retardant.

DIRECTIONS FOR USE: SHAKE WELL BEFORE USING. FOR USE WITH PESTICIDES REGISTERED FOR AGRICULTURAL, AQUATIC, FORESTRY, INDUSTRIAL, MUNICIPAL, NONCROPLAND, ORNAMENTAL, RIGHTS-OF-WAY, TURF AND OTHER APPROPRIATE USES.

The addition of adjuvants to some pesticides and/or fertilizers and their tank mixes may cause phytotoxicity to the foliage and/or fruit of desirable crops. Before using DIRECT, the applicator or crops advisor must have experience with the pesticide combination, including DIRECT, or must have conducted a phytotoxicity test. Excessive use rates of DIRECT will cause a thickened or stringy solution. Dilution with water and at least 1 hour of continuous recirculation will help correct this condition.

USE RATES

As a general guideline, use 1-4 ounces per 100 gallons of spray solution. Always start with the lower dosage and increase as needed to address specific spray conditions. If spray fines are not eliminated, add additional quantities of DIRECT in 1-ounce increments until desired results are achieved.

MIXING

Always conduct a jar test first, to ensure compatibility.

SHAKE WELL BEFORE USING

See special mixing instruction for tank mixes which include Roundup® Ultra and/or wettable powder tank mixes.

General use tank mixes

1. Complete normal mixing and filling procedures.
2. Add DIRECT to the spray solution last at the point of greatest agitation.
3. Continue agitation for at least 3 minutes before spraying and maintain agitation during application.

Wettable powder tank mixes

1. Fill tank one-half full of desired water volume.
2. Add wettable powders
3. Add DIRECT to the spray solution at the point of greatest agitation.
4. Under agitation, add other pesticides and adjuvants in the appropriate order.

Roundup Ultra tank mixes

1. Fill tank one-half full of desired water volume.
2. Add DIRECT to the spray solution at the point of greatest agitation.
3. Under agitation, add other pesticides and adjuvants in the appropriate order.
4. Add Roundup Ultra last.

STORAGE AND DISPOSAL

Store in original container only. If product freezes, bring to room temperature and mix thoroughly before use. Prevent water from being introduced into the contents of this package. Do not reuse empty container and dispose of it in accordance with federal, state and local regulations.

CONDITIONS OF SALE

Note: Read the information contained herein before buying or using this product. If the stated terms are not acceptable, return the product at once, unopened. It is critical that this product be used and mixed only as specified on this label. Neither the manufacturer nor the seller makes any representation or warranty, expressed or implied, with respect to the results from the use of this material. Buyer and user assume all risks of use and/or handling. Precision Laboratories, Inc. warrants that this material is reasonably fit for use as specified on this label. No agent or representative is authorized to make any other representations concerning this material. Unforeseen factors beyond Precision's control prevent elimination of risks in connection with the use of its chemicals. Such risks include, but are not limited to, damage to plants and/or crops to which the material is applied, or lack of complete control and damage caused by drift to other plants or crops. Such risks may occur even though the product is reasonably fit for use as stated herein and even though label directions are followed. Follow directions carefully. Timing, mixture, method of application, weather and other conditions are influencing factors in the use of this product and are beyond the control of the seller. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of the company or any other seller for any and all losses, personal injuries or damages resulting from the use of this product, shall be the purchase price paid by the user or buyer for the quantity of product involved.

S P E C I M E N L A B E L